Intensity variation of gamma radiation on ground level interface in São Jose dos Campos, SP, Brazil

Variação de intensidade da radiação gama na interface ao nível do solo em São José dos Campos, SP, Brasil

DOI: 10.46814/lajdv3n4-065

Recebimento dos originais: 01/05/2021

Aceitação para publicação: 31/06/2021

Alexandre Santos de Souza

Bachelor's Degree in Meteorology from UFPA – Federal University of Pará in 2004. Received his Specialization in Aeronautical Meteorology at ICEA – Airspace Control Institute in 2010. He is currently Captain Specialist in Meteorology at the Ministry of Defense (Command of Aeronautics / Institute of Aeronautics and Space) and Master's student in the Space Sciences and Technologies at the ITA (Technological Institute of Aeronautics), in São José dos Campos, SP, Brazil

E-mail: alexandreass2@fab.mil.br alxsza1973@gmail.com

Cleber Souza Correa

Bachelor's Degree in Meteorology from the Federal University of Pelotas (1987), a Master's in Remote Sensing from the Federal University of Rio Grande do Sul (1997) and a Ph.D. in Water Resources and Environmental Sanitation from the Federal University of Rio Grande do Sul (2005). He was a researcher at the Institute of Airspace Control (ICEA) and is currently a researcher at the Institute of Aeronautics and Space (IAE), acting as a collaborator in the Postgraduate Program in Space Science and Technologies (PG/CTE-E) of the Technological Institute of Aeronautics (ITA).

As an Officer in the Air Force Command, he has a Specialization course in Aeronautical Meteorology at ICEA in 1988 and the Course of Command and General Staff of Aeronautics at the University of the Air Force (UNIFA) in Rio de Janeiro, held in 2012. He has experience in the area of Geosciences, with an emphasis on Meteorology, acting on the following topics: Micrometeorology, Mesoscale Meteorology, Climatology, Numerical Modeling and Statistics.

E-mail: clebercsc@fab.mil.br cleber.voy@gmail.com

Inacio Malmonge Martin

Bs degree in Physics from UNESP – State University of São Paulo in 1967. Received his Msdegree in Physics at ITA (Technological Institute of Aeronautics) in 1968. Received his Docteurd'Specialitéat Université Toulouse III (Paul Sabatier) in Toulouse, France in 1970. Received his Docteurd'Etat in 1974 from Universitéof Toulouse III (Paul Sabatier) in Toulouse,France.He is currently associate professor (graduate and pós-graduate students) in the Physics Department at the ITA, in São José dos Campos, SP, Brazil

E-mail: martin@ita.br inaciomalmonge@uol.com.br

ABSTRACT

In the period June 22 to September 24, 2012 São José dos Campos region experienced dry climate with low relative humidity. The net rain intensity measured in this period was 60 mm/h and the relative humidity range between 30% to 40% on most days. The intensity of gamma radiation from 0.03 to 10.0 MeV, in the measurement period presents a daily fluctuation well clearly, and some small increases when there is rainfall. These measures were realized at ITA's campus with a scintillator of NaI(Tl), photomultiplier and associated electronics laptop to a Dell 630 PC. A data acquisition interface purchased from (Aware Eletronics, USA) provides a file with measurements versus time, in the range

from minute to minute recorded in txt. It was observed that the variation of gamma radiation in the interface soil/air at the region is originating from the local radon gas dynamics and of the presence of wind shear near the ground level.

Keywords: Atmosphere, Radon gas, Environmental gamma radiation.

RESUMO

No período de 22 de junho a 24 de setembro de 2012, a região de São José dos Campos experimentou um clima seco com baixa umidade relativa. A intensidade líquida de chuva medida neste período foi de 60 mm/h e a umidade relativa varia entre 30% a 40% na maioria dos dias. A intensidade de radiação gama de 0,03 a 10,0 MeV, no período de medição apresenta uma flutuação diária bem clara, e alguns pequenos aumentos quando há chuva. Estas medidas foram realizadas no campus da ITA com um cintilador de NaI(Tl), fotomultiplicador e laptop eletrônico associado a um PC Dell 630. Uma interface de aquisição de dados adquirida da (Aware Eletronics, EUA) fornece um arquivo com medidas versus tempo, na faixa de minutos a minutos gravados em txt. Foi observado que a variação da radiação gama na interface solo/ar na região é originada pela dinâmica local do gás radônio e da presença de cisalhamento do vento próximo ao nível do solo.

Palavras-chave: Atmosfera, gás radon, radiação gama ambiental.

1 INTRODUCTION

It is common knowledge that we live in an environment which comprises ionizing radiation (X-rays, gamma rays). These radiations [1], in normal situations, are produced by natural phenomena such as cosmic rays from different sources and the decay of radioactive elements present in the planet's atmosphere and crust. Ionizing radiation can also be produced by human activities such as those related to nuclear power generation and use in medicine. Although ionizing radiations are part of our daily lives, few students or researchers have the opportunity to perform measurements and observations of the background radiation. To overcome this deficiency in education we started a pilot project comprising measures and observations of environmental radiation background with students of the Technological Institute of Aeronautics (ITA, SP) and the University of Taubate (UNITAU, SP) using compact and portable meters to observe gamma radiation.

The geological radiation source is largely the result of the decay of radioactive isotopes of potassium (40K), uranium (238U) and thorium (232Th) [2]. In radioactive decay, nuclei may emit alpha radiation, beta, gamma or X-rays, in addition, besides, are formed the natural radioactive isotopes belonging to each series decay aforementioned. The 235U has a half-life of 713 million years whereas the 238U has a half-life of 4,5 billion year. For this reason there is a much smaller amount of 235U compared to 238U [3].

Radon gas (222Rn) is the other important source of ionizing radiation on interface soil/air which likewise is formed by uranium and thorium decay. In turn, the radioactive decay of radon produces

^{218}Po , ^{214}Pb and ^{214}Bi , These isotopes may be measured in the air [4-7]. A relevant fact related to radon is that during the rainfall its soil concentration may rise due to the transport of that gas through raindrops. The secondary cosmic radiation produced on lower atmosphere has a little variation with time; however, it varies accordingly to latitude and altitude on the earth's surface [8]. Secondary cosmic radiation produces "extensive showers" of energetic particles and photons which arrives to earth's surface [9]. Artificial ionizing radiations are those man-made in several kinds of activities such as medicine, research, and industry. Nevertheless, these radiations source are confined and over control. In Brazil have not yet been made the measures on the territory, of radiation from soil, produced by atmospheric radon, cosmic origin and artificial. But in other countries these values are known, for example in Portugal 56% of ground level radiation is produced by atmospheric radon, 8% produced by soil, 8% is cosmic origin, 17% is artificially produced, and 11% by other sources [10].

2 MATERIALS AND METHODS

Gamma radiation measurements was realized with a crystal scintillator of NaI(Tl) mounted in an aluminum cell and connected to a photomultiplier tube (Aware Electronics, USA). This detector was coupled to a PC for storing the data series. The scintillator, coupled to the photomultiplier and associated electronics, was previously calibrated with standard radioactive sources in a laboratory.

Figure 1 - Portable system for measurements used at this work.

The background radiation measurement is collected each minute uninterruptedly during one day or until three months. The methods used to study the data sets are statistics and Fourier analysis. The portable system to measure the environment gamma radiation which was used in this work is shown in Fig.1.

3 RESULTS AND DISCUSSIONS

Measurements of gamma radiation were performed at the Physics Department of the ITA in São José dos Campos, SP, from 22 June 2012 to 24 September 2012 at fixed counting intervals of 1 minute continuously throughout the period. In Fig. 2 shows the radiation collected by the scintillator per minute versus time of measurement. Note that the measurements of X-ray photons and gamma occur so omnidirectional, irrespective of direction. During this period the atmosphere near the Earth's surface was extremely dry with low humidity in the afternoons, and around 30% in most parts of the days analyzed. The rains were all weak and of low intensity (mm/h) distributed among 10/07 and in 07/17, as shown in Fig. 2, a total of 54 mm/h [11]. From the day 07/17 until day 09/21, the local atmosphere was very dry. On 09/22 the first rain arrived with very low intensity, near 6 mm/h. Even with low intensity of precipitation in those days it was noted a correlation with the intensity of gamma radiation observed.

Figure 2 - Measurements of gamma radiation and precipitation [11] in São José dos Campos the period from 22 June 2012 to 24 September 2012.

Fig. 3 shows the variation between the above-mentioned measure relative humidity experienced in the region. For each day the average value was determined that moisture. The data used to calculate the average value of relative humidity were obtained through a datalogger system [12].

In every period measures the radiation varies with a cycle of 24 hours perfectly shown in the graph of Fig. 4 (power spectra FFT). This cycle is quite evident in measurements with dry weather interference where rain is almost null. The radiation in this period without rain always increases between dawn until the sun 11-12 hours local, this is due to greater presence of radon gas (^{222}Rn) which decays after 3.8 days in ^{214}Pb and ^{214}Bi , emitting gamma photons belonging to the energy range of the scintillator measuring. At the time of rain radon gas also increases due to the phenomenon of washing the lower atmosphere where the radon gas is diluted in suspended water droplets.

Figure 4 - The power spectra (FFT) obtained for the period.

4 CONCLUSION

The measurements of gamma radiation from 0.03 to 10.0 MeV were performed continuously from 06/22 to 09/24, 2012, per each minute on the campus of the Physics Department of the ITA in São Jose dos Campos, SP, Brazil in ground level interface. Atmospheric pressure, relative humidity and temperature were also measured at the same time and location with the same time interval of minutes. The analysis of these measurements clearly shows how the radon gas provokes influence of the dry period, causing a well-defined cycle of 1 day. The average relative humidity during all days clearly shows its decline from the month of June to September 2012 where occurred the first small rainfall of this period.

ACKNOWLEDGEMENTS

To Fortunato Rezende Guimarães financing instruments, to CNPq process projects 480407/2011-8 and 305145/2009-6, to UNITAU and ITA for the logistical and technical support.

REFERENCES

- Eisenbud, M. and Gesell, T. F., 1997. Environmental Radioactivity from Natural, Industrial & Military Sources, 4th ed. London: Academic Press.
- Buivan, N.A., Martin, I.M. and Turtelli, A. Jr., 1988 Measurements of natural radioactivity at different atmospheric depths. Revista Geofísica, IPGH, numero 28, México.
- Magill, J. and Galy, J., 2004. Radioactivity Radionuclides Radiation, New York: Springer.
- Hosler, J C. R., 1966. "Meteorological effects on atmospheric concentration of radon (Rn^{222}), RaB (Pb^{214}), and RaC (Bi^{214}) near the ground," Monthly Weather Review, vol. 94, no.2, pp 89-99,
- Pearson, J. E. and Jones, G. E., 1966. "Soil concentrations of "emanating radium-226" and the emanation of radon-222 from soils and plants," Tellus XVIII, vol. 2, pp. 656-662.
- Fujinami, N., 2009. "Study of radon progeny distribution and radiation dose rate in the atmosphere," Jpn. J. Health Phys., vol. 44, no. 1 pp. 88-94.
- Ichiji, T. and Hattori, T. "Continuous measurement of environmental gamma radiation in Tokyo using Ge semiconductor detector," IRPA homepage.[Online].Available:<http://www.irpa.net/irpa10/cdrom/00611.pdf>.
- Martin, I. M., 29 sept. 1971. Variation du Flux des Neutrons et des Rayons Gamma d'origine cosmique en fonction de la latitude, These de Doctorate de Spécialité, Université de Toulouse, France.
- Grieder, P. K. F. Ed., 2001. Cosmic Rays at Earth, Amsterdam: Elsevier.
- Lima, A. A. M, 2007 – MSc Thesis – University of Coimbra, Portugal.
- Wireless rain meter Oregon scientific RGR122, TFA cod. N. KW-9015 <http://foro.tiempo.com/pluviometro-oregon-scientific-rgr122-t9719.0.html> 10/10/2012.
- Martin, I.M., JG Silvério, T. Mitsuo; system datalogger-arduino measures in real time and available on the internet; [http://200.145.13.145:99/datalogger/\(10/10/2012\)](http://200.145.13.145:99/datalogger/(10/10/2012)).